

- Consumer Services, Division of Plant Industry.
- Knolhoff LM, Heckel DG, 2014. Behavioral assays for studies of host plant choice and adaptation in herbivorous insects. *Annual Review of Entomology*, 59: 263–278.
- Köllner TG, Held M, Lenk C, Hiltbold I, Turlings TC, Gershenzon J, Degenhardt J, 2008. A maize (*E*)- $\beta$ -caryophyllene synthase implicated in indirect defense responses against herbivores is not expressed in most American maize varieties. *The Plant Cell*, 20(2): 482–494.
- König G, Brunda M, Puxbaum H, Hewitt CN, Duckham SC, Rudolph J, 1995. Relative contribution of oxygenated hydrocarbons to the total biogenic VOC emissions of selected mid-European agricultural and natural plant species. *Atmospheric Environment*, 29(8): 861–874.
- Li CR, Xia WS, Wang FL, 2007. First record of *corythucha ciliata* (Say)(Hemiptera; Tingidae), in China. *Acta Zootax. Sin.*, 32 (4): 944–946. [李传仁, 夏文胜, 王福莲, 2007. 悬铃木方翅网蝽在中国的首次发现. *动物分类学报*, 32(4): 944–946.]
- Maceljski M, Balarin I, 1972. Preliminary note on the appearance of a new species of insect pest in Yugoslavia—the bug *Corythucha ciliata* (Say) (Tingidae, Heteroptera). *Acta Entomologica Jugoslavica*, 8(1/2): 105–106.
- Matsuki M, Foley WJ, Floyd RB, 2011. Role of volatile and non-volatile plant secondary metabolites in host tree selection by Christmas beetles. *Journal of Chemical Ecology*, 37(3): 286–300.
- Tholl D, Sohrabi R, Huh JH, Lee S, 2011. The biochemistry of homoterpenes—common constituents of floral and herbivore-induced plant volatile bouquets. *Phytochemistry*, 72(13): 1635–1646.
- Turlings TC, Tumlinson JH, Lewis WJ, 1990. Exploitation of herbivore-induced plant odors by host-seeking parasitic wasps. *Science*, 250(4985): 1251–1253.
- Wang FL, Li CR, Liu WX, Wan FH, 2008. Advance in biological characteristics and control techniques of the new invasive sycamore lacebug (*Corythucha ciliata*). *Scientia Silvae Sinicae*, 44(6): 137–142. [王福莲, 李传仁, 刘万学, 万方浩, 2008. 新入侵物种悬铃木方翅网蝽的生物学特性与防治技术研究进展. *林业科学*, 44(6): 137–142.]

\*\*\*\*\*



## 马力筋红天牛 *Tetraopes femoratus* LeConte

马力筋红天牛为鞘翅目 Coleoptera 天牛科 Cerambycidae 昆虫, 主要分布在北美洲, 美国中西部及墨西哥中部常见, 取食各种马利筋属 *Asclepias* 植物, 如乳草 *A. fascicularis*, 沼泽乳草 *A. incarnata* 等。成虫 3 cm 左右, 体色鲜红, 触角黑色节间白色, 前胸背板中部及两侧有大的突起, 鞘翅红色, 有 1~2 对小的黑色斑点。封面照片 2015 年 9 月 14 日拍摄于美国亚利桑那州图森市 (Tucson, Arizona) 的一个峡谷 Madera Canyon。

(姜春燕, 中国科学院动物研究所)