

Seven new species of jumping spiders (Araneae, Salticidae) from Xishuangbanna, China

Cheng Wang¹, Shuqiang Li²

1 College of Agriculture and Forestry Engineering and Planning, Tongren University, Tongren, Guizhou, 554300, China **2** Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China

Corresponding author: Shuqiang Li (lisq@ioz.ac.cn)

Academic editor: Yuri Marusik | Received 3 June 2020 | Accepted 23 August 2020 | Published 16 September 2020

<http://zoobank.org/90FD0800-76BF-488D-AC46-EF93E4BCCE21>

Citation: Wang C, Li S (2020) Seven new species of jumping spiders (Araneae, Salticidae) from Xishuangbanna, China. ZooKeys 968: 43–69. <https://doi.org/10.3897/zookeys.968.55047>

Abstract

Seven new species of jumping spiders collected from Xishuangbanna Tropical Botanical Garden in Yunnan, China, are diagnosed and described: *Charippus yinae* **sp. nov.** (♂♀), *Chinattus inflatus* **sp. nov.** (♂), *Indomarengo yui* **sp. nov.** (♂), *Phintella banna* **sp. nov.** (♂♀), *P. mii* **sp. nov.** (♂♀), *Simaetha menglun* **sp. nov.** (♂♀) and *S. pengi* **sp. nov.** (♂♀). *Charippus yinae* **sp. nov.** is the second species of the genus *Charippus* Thorell, 1895, which was previously known only from one sex.

Keywords

distribution, genus, salticid, taxonomy, Yunnan

Introduction

Xishuangbanna is a key biogeographic area and a biodiversity hotspot in China (Myers 1988). Spider diversity in the area is high, with 782 species spanning 305 genera in 46 families (Li 2020). The Salticidae Blackwall, 1841 from this region have been studied by Peng and Yin (1991), Song (1991), Peng (1995), Xie and Peng (1995), Peng and Kim (1997), Song and Zhu (1998), Xiao (2002), Xiao and Wang (2004), Cao et al. (2016), Wang and Li (2020), and Lin and Li (2020). These studies have resulted

in the description of 45 new species and increased the total salticid species number to 121 in the last 30 years (Lin and Li 2020). At least 19 salticid species described as new from Xishuangbanna are known only from a single sex, and sparsely collected new species are frequently being discovered as a result of the ongoing project “All Species Inventory” of spiders from the Xishuangbanna Tropical Botanical Garden. Our results thus far indicate that the salticid fauna of Xishuangbanna remains insufficiently known and requires further research.

A taxonomic study on the recently available salticid samples of Xishuangbanna has revealed seven species that are new to science and are described here. These results increase the total number of salticid species in the region to 128.

Materials and methods

Specimens were collected by fogging and sieving leaf litter in the tropical rainforest of Xishuangbanna Tropical Botanical Garden, Yunnan, China. All specimens were preserved in 75% ethanol and are deposited in the Institute of Zoology, Chinese Academy of Sciences (IZCAS) in Beijing, China. The specimens were examined with an Olympus SZ51 stereomicroscope. After dissection, the epigyne was cleared in trypsin enzyme solution before examination and imaging. Images of the copulatory organs and habitus were taken with a Kuy Nice CCD mounted on an Olympus BX51 compound microscope. Compound focus images were generated using Helicon Focus v. 6.7.1.

All measurements are given in millimeters. Leg measurements are given as: total length (femur, patella + tibia, metatarsus, tarsus). References to figures in the cited papers are listed in lowercase type (fig. or figs); figures in this paper are noted with an initial capital (Fig. or Figs). Abbreviations used in the text and figures are as follows:

AAM	anterior atrial margin;	FD	fertilization duct;
AER	anterior eye row;	H	hood;
AERW	anterior eye row width;	LAM	lateral atrial margin;
ALE	anterior lateral eye;	LP	lamellar process;
AME	anterior median eye;	PERW	posterior eye row width;
AS	anterior chamber of spermatheca;	PL	posterior lobe;
BP	basal plate;	PLE	posterior lateral eye;
CD	copulatory duct;	PS	posterior chamber of spermatheca;
CO	copulatory opening;	RFA	retrolateral femoral apophysis;
CP	cymbial process;	RTA	retrolateral tibial apophysis;
E	embolus;	S	spermatheca;
EC	embolic coil;	SD	sperm duct;
EFL	eye field length;	TB	tegular bump.
F	fold;		

Taxonomy

Family Salticidae Blackwall, 1841

Genus *Charippus* Thorell, 1895

Type species. *Charippus errans* Thorell, 1895

Comments. This monotypic genus is only known from the descriptions of the generotype male (Thorell 1895; WSC 2020). *Charippus* is similar to *Cytaea* Keyserling, 1882 by the general shape of the copulatory organs, especially the spiral embolus and the long and coiled copulatory ducts but can be distinguished by the following: 1) the carapace is slightly narrowed anteriorly vs almost square in *Cytaea* species; 2) the color of the habitus is dark rather than light, as in *Cytaea* species, indicating that *Charippus* likely comprises leaf-litter dwellers rather than foliage dwellers, as in *Cytaea*; 3) the RTA is distinctly longer than in *Cytaea* species; 4) the male chelicerae only have two promarginal teeth vs at least three promarginal teeth in *Cytaea* species; 5) the epigyne lacks a distinct median septum, which is present in *Cytaea* species.

Charippus yinae sp. nov.

<http://zoobank.org/97A61C13-E1F2-49D7-8074-0404C15CDD05>

Figs 1, 2

Type material. *Holotype* ♂ (IZCAS-Ar40601), CHINA: Yunnan: Xishuangbanna, Mengla County, Menglun Town, Menglun Nature Reserve, Xishuangbanna Tropical Botanical Garden, Leprosy Village (21°53.59'N, 101°17.30'E, ca 550 m), 29.04.2019, Y. Tong et al. leg. *Paratypes* 1 ♀ (IZCAS-Ar40602), same data as holotype; 1 ♀ (IZCAS-Ar40603), Flower Garden (21°55.80'N, 101°45.41'E, ca 530 m), 1.08.2018, C. Wang et al. leg.

Etymology. The species name is a patronym in honor of the late Professor Changmin Yin, one of the pioneers of spider taxonomy of China; noun (name) in genitive case.

Diagnosis. The male of *Charippus yinae* sp. nov. resembles *C. errans* Thorell, 1895 by the habitus, shape of the palp, and the cheliceral dentition, but can be easily distinguished by the embolus originating at 9:00 o'clock rather than approximately 11:30 o'clock as in *C. errans* (Wanless 1988: fig. 40F), and the RTA is curved towards the bulb medially in retrolateral view, instead of curved towards the bulb terminally as in *C. errans* (Wanless 1988: fig. 40H). The female resembles *Cytaea alburna* Keyserling, 1882 by the very long, coiled copulatory ducts and the small spermathecae but can be easily distinguished by the following: 1) the carapace is slightly narrowed anteriorly rather than almost square as in *C. alburna* (Davies and Žabka 1989: 226); 2) the median septum is absent vs present in *C. alburna* (Davies and Žabka 1989: 226); 3) the chelicerae have only two promarginal teeth vs three promarginal teeth as in *C. alburna* (Davies and Žabka 1989: 226).




Figure 1. Male palp of the holotype of *Charippus yinae* sp. nov. **A** prolateral **B** retrolateral **C** ventral **D** dorsal. Scale bars: 0.1.




Figure 2. *Charippus yinae* sp. nov., female paratype and male holotype **A** epigyne, ventral **B** vulva, dorsal **C** holotype habitus, dorsal **D** holotype habitus, ventral **E** paratype habitus, dorsal **F** holotype carapace, frontal **G** holotype chelicerae, dorsal. Scale bars: 0.1 (**A**, **B**, **G**); 0.5 (**C**-**F**).

Description. Male (Figs 1, 2C, D, F, G). Total length 3.91. Carapace 1.98 long, 1.56 wide. Abdomen 2.04 long, 1.48 wide. Clypeus 0.10 high. Eye sizes and inter-distances: AME 0.38, ALE 0.22, PLE 0.20, AERW 1.28, PERW 1.24, EFL 0.97. Legs: I 3.83 (1.15, 1.61, 0.68, 0.39), II 3.53 (1.07, 1.39, 0.68, 0.39), III 3.78 (1.22, 1.29, 0.88, 0.39), IV 4.27 (1.22, 1.49, 1.07, 0.49). Carapace dark brown, with an irregular red area in the middle of the thoracic part, bearing dense white setae on the cheeks and white and golden setae dorsally. Clypeus dark, with sparse white setae. Fovea longitudinal, dark. Chelicerae red to dark brown, with two promarginal teeth and one retro-marginal fissident with two cusps. Endites red-brown, paler inner margin. Labium dark brown. Sternum somewhat shield-shaped, with sparse setae. Legs I red to dark brown; other legs yellow, with dark markings. Abdomen suboval, dorsum hirsute, with two pairs of faint pale patches anteriorly and a pair of posterolateral falcate bands followed by several chevrons; venter dark anteriorly, covered by small spots and thin brown setae.

Palp (Fig. 1A–D): tibia wider than long in ventral view, with a RTA approximately two times the tibial length, slightly curved medially and blunt at the tip in retrolateral view; cymbium hirsute, proximodorsally with a small process (Fig. 1D, CP) near the RTA base; bulb swollen, with sperm duct strongly curved retrolaterally; embolus originating at around 9:00 o'clock, coiled 1.5 times.

Female (Fig. 2A, B, E). Total length 3.83. Carapace 1.83 long, 1.38 wide. Abdomen 2.13 long, 1.45 wide. Clypeus 0.10 high. Eye sizes and inter-distances: AME 0.36, ALE 0.23, PLE 0.21, AERW 1.20, PERW 1.14, EFL 0.88. Legs: I 3.08 (0.98, 1.27, 0.49, 0.34), II 2.88 (0.95, 1.10, 0.49, 0.34), III 3.32 (1.10, 1.15, 0.73, 0.34), IV 3.58 (1.12, 1.29, 0.83, 0.34). Habitus similar to that of male except slightly paler.

Epigyne (Fig. 2A, B): wider than long, with a pair of arc-shaped anterior atrial margins and a pair of lateral atrial margins; copulatory openings small, slit-shaped, located anterolaterally; copulatory ducts very long, curved into a U-shape at the origin, with anterior tubercles and complex coils; spermathecae suboval, separated from each other by almost two times their width; fertilization ducts nearly cordiform.

Distribution. Known only from the type locality in Yunnan, China.

Genus *Chinattus* Simon, 1876

Type species. *Habrocestoides szechwanensis* Prószyński, 1992

Comments. The genus is represented by 17 nominal species restricted to Asia, except *C. parvulus* (Banks, 1895), which is known from the USA and Canada. Diagnostic drawings have been made for all species in the genus, and nearly all have limited distributional ranges: 11 species are endemic, and five species are known only from two countries. Six species are known only from a single sex: two from males and four from females (all from China). To date, 12 species are recorded from China. Of those, eight are endemic (WSC 2020).

***Chinattus inflatus* sp. nov.**

<http://zoobank.org/6BD4F191-4595-492A-8794-6480493017BB>

Figs 3, 4

Type material. *Holotype* ♂ (IZCAS-Ar40604), CHINA: Yunnan: Xishuangbanna, Mengla County, Menglun Town, Menglun Nature Reserve, Xishuangbanna Tropical Botanical Garden, tropical rainforest (21°55.20'N, 101°16.21'E, ca 550 m), 30.04.2019, Y. Tong et al. leg. *Paratype* 1♂ (IZCAS-Ar40605), same data as holotype.

Etymology. The specific name refers to the inflated femur of the male palp; adjective.

Diagnosis. *Chinattus inflatus* sp. nov. resembles *C. wengnanensis* Cao & Li, 2016 in the general shape of the habitus and male palp but can be easily distinguished by the inflated palpal femur, the femoral apophysis and the long embolus. The new species also resembles species of the genus *Grayenulla* Zabka, 1992 by the inflated palpal femur and the femoral apophysis but can be distinguished by leg III, which is one of the shortest legs rather than distinctly the longest as in species of *Grayenulla*. Additionally, the clypeus lacks bristles, whereas there are three central bristles in species of *Grayenulla* (Zabka 1992).

Description. Male (Figs 3, 4). Total length 2.72. Carapace 1.53 long, 1.14 wide. Abdomen 1.19 long, 0.92 wide. Clypeus 0.10 high. Eye sizes and inter-distances: AME 0.36, ALE 0.21, PLE 0.17, AERW 1.13, PERW 1.08, EFL 0.57. Legs: I 2.82 (0.85, 1.12, 0.46, 0.39), II 2.38 (0.73, 0.85, 0.41, 0.39), III 2.65 (0.85, 0.85, 0.56, 0.39), IV 2.75 (0.88, 0.85, 0.63, 0.39). Carapace brown, dark in eye field, bearing white, thin setae on the lateral submargin, and golden, thin setae on the cheeks and cephalic region. Clypeus dark. Fovea longitudinal, short, bar-shaped. Chelicerae red-brown, with two promarginal teeth and one retromarginal fissident. Endites, labium and sternum paler than chelicerae. Legs red-brown to dark brown, with pale rings on femora and metatarsi. Abdomen suboval, dorsum brown, speckled, with three pairs of irregular white spots and two pairs of muscle depressions; venter darker than dorsum, with a pair of off-white longitudinal bands laterally and a pair of dotted lines medially.

Palp (Fig. 3A–D): femur inflated, more than 1.5 times longer than wide, with a subtriangular retrolateral apophysis proximally; tibia wider than long, with a sclerotized, short RTA extending towards the bulb in retrolateral view and almost completely hidden by the bulb in ventral view; posterior lobe well-developed, with oval margin; embolus flat and straight, tapered to a pointed tip.

Female. Unknown.

Distribution. Known only from the type locality in Yunnan, China.

Comments. The species is placed into this genus because it generally resembles *C. wengnanensis* Cao & Li, 2016. It is described only based on males, and so there is a possibility that it is conspecific to one of the congeners known only from females.




Figure 3. Male palp of the holotype of *Chinattus inflatus* sp. nov. **A** prolateral **B** retrolateral **C** ventral **D** bulb, ventral. Scale bars: 0.1.



Figure 4. *Chinattus inflatus* sp. nov., male holotype **A** habitus, dorsal **B** habitus, ventral **C** carapace, frontal **D** chelicerae, dorsal. Scale bars: 0.5 (**A–C**); 0.1 (**D**).

Genus *Indomarengo* Benjamin, 2004

Type species. *Indomarengo sarawakensis* Benjamin, 2004

Comments. This genus only contains four species, all endemic: one from Borneo, two from Indonesia and one from India. Two are known only from a single sex: *I. chandra* Benjamin, 2004 from males and *I. thomsoni* (Wanless 1978) from females (WSC 2020). The genus can be distinguished from all other Ballini genera, except *Sadies* Wanless, 1984, *Leikung* Benjamin, 2004, and *Afromarengo* Benjamin, 2004, by the presence of a carapace protuberance. It can be distinguished from *Afromarengo* and *Leikung* by the short embolus, which has less than two spirals, and from *Sadies* by the lack of leaf-like carapace scales (Benjamin 2004).

Indomarengo yui sp. nov.

<http://zoobank.org/6CA2DCC9-71E9-488B-9A3F-29B3E59E09AC>

Figs 5, 6

Type material. *Holotype* ♂ (IZCAS-Ar40606), CHINA: Yunnan: Xishuangbanna, Mengla County, Menglun Town, Menglun Nature Reserve, garbage dump, secondary tropical rainforest (21°54.30'N, 101°16.78'E, ca 620 m), 26.04.2019, H. Yu et al. leg. *Paratype* 1♂ (IZCAS-Ar40607), same data as holotype.

Etymology. The specific name is a patronym after Prof. Hao Yu (Guiyang, China), one of the collectors of the new species; noun (name) in genitive case.




Figure 5. Male palp of the holotype of *Indomarengo yui* sp. nov. **A** prolateral **B** retrolateral **C** ventral **D** bulb, retrolateral. Scale bars: 0.1.



Figure 6. *Indomarengo yui* sp. nov., male holotype **A** habitus, dorsal **B** habitus, lateral **C** habitus, ventral **D** chelicerae, dorsal **E** left leg I, prolateral. Scale bars: 0.5 (**A–C, E**); 0.1 (**D**).

Diagnosis. *Indomarengo yui* sp. nov. resembles *I. chavarapater* Malamel, Prajapati, Sudhikumar & Sebastian, 2019 by the general shape of the palp but can be distinguished by the RTA, which is distally curved towards the bulb in retrolateral view vs almost straight and directed anteriorly in *I. chavarapater* (Malamel et al. 2019: figs 11, 15); and by the tibia, which is wider than long vs almost as long as wide in *I. chavarapater* (Malamel et al. 2019: figs 10, 11, 15, 16).

Description. Male (Figs 5, 6). Total length 3.12. Carapace 1.36 long, 0.85 wide. Abdomen 1.67 long, 0.76 wide. Clypeus 0.05 high. Eye sizes and inter-distances: AME 0.29, ALE 0.12, PLE 0.12, AERW 0.75, PERW 0.78, EFL 0.53. Legs: I 3.46 (0.93, 1.51, 0.80, 0.22), II 1.79 (0.54, 0.66, 0.37, 0.22), III 1.67 (0.49, 0.59, 0.37, 0.22), IV 2.09 (0.63, 0.78, 0.46, 0.22). Carapace flat, red-brown to dark brown, there are white setae behind the PLEs and posteriorly, with a distinct protuberance on the thoracic region. Clypeus dark. Fovea indistinct. Chelicerae red-brown, with three retromarginal teeth and two promarginal teeth. Endites and labium dark brown. Sternum red-brown, elongated and tapered posteriorly, covered with thin setae. Legs I enlarged, with inflated tibia bearing ventral leaf-shaped scales and five spines; other legs pale, with dark stripes laterally on femora and patellae. Abdomen elongated oval, dorsum brown to dark brown, constricted at first anterior third, with white setae at lateral margins; venter pale to brown.

Palp (Fig. 5A–D): tibia wider than long, with a thin RTA approximately two times the tibial length, widened at the base and curved towards bulb terminally; bulb inflated, divided by a furrow; embolus short, coiled, forming two circles, and partly covered by a membranous structure.

Female. Unknown.

Distribution. Known only from the type locality in Yunnan, China.

Comments. The species has a distinct carapace protuberance, which is only known to occur in species of the genera *Indomarengo* Benjamin, 2004, *Sadies* Wanless, 1984, *Leikung* Benjamin, 2004, and *Afromarengo* Benjamin, 2004 of the Ballini. Based on similarities such as the short embolus with less than two spirals and lack of leaf-like carapace scales (more than two spirals in *Leikung* and *Afromarengo* species and leaf-like carapace scales present in *Sadies* species) in addition to closely resembling *I. chavarapater* Malamel, Prajapati, Sudhikumar & Sebastian, 2019 by the habitus and the shape of the male palp, we place the spider into this genus.

Genus *Phintella* Strand, 1906

Type species. *Telamonia bifurcilinea* Bösenberg & Strand, 1906

Comments. *Phintella* is a rather large genus, represented by a group of small and colorful spiders, typically covered with metallic iridescent scales (Luong et al. 2016). A total of 60 nominal species have been described, primarily from the Afrotropical, Oriental, and Palearctic regions. Diagnostic drawings have been made for all species in the genus, and 27 species are known only from a single sex: 13 from females and 14 from males. To date, 47 species are known from Asia, including 29 from China. Among the Chinese *Phintella*, 16 are endemic and 12 are known from only a single sex: six from males and six from females (WSC 2020). Based on the continuous discovery of undescribed species in our recent fieldwork, it is likely that the true diversity of Chinese *Phintella* is much greater than currently known.

Phintella banna sp. nov.

<http://zoobank.org/C3900F5F-68F0-48E1-805F-53069417C171>

Figs 7, 8

Type material. *Holotype* ♂ (IZCAS-Ar40608), CHINA: Yunnan: Xishuangbanna, Mengla County, Menglun Town, Menglun Nature Reserve, Leprosy Village (21°53.59'N, 101°17.30'E, ca 550 m), 4.05.2019, Y. Tong et al. leg. *Paratypes* 5♀3♂ (IZCAS-Ar40609–40616), same data as holotype; 1♀3♂ (IZCAS-Ar40617–40620), Xishuangbanna Tropical Botanical Garden, tropical rainforest (21°55.35'N, 101°16.36'E, ca 610 m), 7.08.2018, C. Wang et al. leg.

Etymology. The species name is derived from the name of the type locality; noun in apposition.

Diagnosis. *Phintella banna* sp. nov. resembles *P. bifurcilinea* (Bösenberg & Strand, 1906) by the small size and the shape of the copulatory organs but can be distinguished

by the following: 1) the lamellar process is visible in ventral view vs obscured in *P. bifurcilinea* (Žabka 1985: fig. 403); 2) the embolus is apically directed anteriorly in ventral view vs directed retrolaterally in *P. bifurcilinea* (Žabka 1985: fig. 403); 3) the chelicerae of the male are underdeveloped, the ratio of the length of the fang to the width of the paturon is about 1:1 vs well-developed chelicerae, with a ratio of almost 2.5:1 in *P. bifurcilinea* (Žabka 1985: fig. 406); 4) the copulatory openings are anterior to the spermathecae vs lateral to the spermathecae in *P. bifurcilinea* (Peng et al. 1993: fig. 527); 5) the copulatory ducts are approximately one-third of the spermathecal diameter vs less than one-fifth of the spermathecal diameter in *P. bifurcilinea* (Peng et al. 1993: fig. 527).

Description. Male (Figs 7, 8C, D, F, G). Total length 2.41. Carapace 1.34 long, 1.02 wide. Abdomen 1.12 long, 0.88 wide. Clypeus 0.10 high. Eye sizes and inter-distances: AME 0.33, ALE 0.17, PLE 0.15, AERW 1.01, PERW 0.98, EFL 0.64. Legs: I 2.73 (0.82, 0.98, 0.59, 0.34), II 2.66 (0.80, 0.93, 0.59, 0.34), III 3.25 (1.01, 1.07, 0.83, 0.34), IV 3.46 (1.10, 1.12, 0.90, 0.34). Carapace dark brown, slightly paler medially in the thoracic area, with yellow and pale scales on the cheeks. Clypeus dark brown. Fovea indistinct. Chelicerae yellow-brown to brown, with one retromarginal tooth and two promarginal teeth. Endites green-brown, pale yellow along the inner edge. Labium wider than long, paler terminally. Sternum green-brown, covered with thin setae. Legs pale to green-brown. Abdomen oval, dorsum hairy, with two pairs of muscle depressions and a transverse white band of scales in the center, followed by recurved dotted lines; venter colored as dorsum, darker medially.

Palp (Fig. 7A–D): tibia wider than long, with a tapered RTA terminally curved towards the bulb and pointed apically in retrolateral view; bulb flat, with a lamellar process near the base of the embolus, approximately three times longer than wide, with a small subtriangular bump and a gently curved posterior lobe reaching the middle of the tibia in ventral view; sperm duct long, strongly curved anteriorly; embolus bar-shaped, almost equal to the length of the lamellar process.

Female (Fig. 8A, B, E). Total length 3.07. Carapace 1.29 long, 1.04 wide. Abdomen 1.76 long, 1.33 wide. Clypeus 0.10 high. Eye sizes and inter-distances: AME 0.32, ALE 0.18, PLE 0.16, AERW 1.04, PERW 1.01, EFL 0.67. Legs: I 2.31 (0.73, 0.80, 0.49, 0.29), II 2.25 (0.71, 0.76, 0.44, 0.34), III 2.88 (0.88, 0.93, 0.73, 0.34), IV 3.31 (1.02, 1.07, 0.88, 0.34). Habitus similar to that of the male except the transverse abdominal white band is indistinct laterally, and the dorsum of the abdomen with a brown, thin, longitudinal band in the middle.

Epigyne (Fig. 8A, B): wider than long, with an arc-shaped basal plate; copulatory openings anteriorly located, slit-shaped, the distance between them almost equal to the spermathecal diameter; copulatory ducts curved anteriorly, and then descend along the longitudinal axis to connect to the postero-ento-lateral part of the spermathecae; spermathecae spherical, separated from each other by less than one-third their diameter; fertilization ducts well-developed, lamellar, anterior to spermathecae.

Distribution. Known only from the type locality in Yunnan, China.




Figure 7. Male palp of the holotype of *Phintella banna* sp. nov. **A** prolateral **B** retrolateral **C** ventral **D** dorsal. Scale bars: 0.1.




Figure 8. *Phintella banna* sp. nov., female paratype and male holotype **A** epigyne, ventral **B** vulva, dorsal **C** holotype habitus, dorsal **D** holotype habitus, ventral **E** paratype habitus, dorsal **F** holotype carapace, frontal **G** holotype chelicerae, dorsal. Scale bars: 0.1 (**A**, **B**, **G**); 0.5 (**C**-**F**).

***Phintella mii* sp. nov.**

<http://zoobank.org/58C57CE9-D503-4CAF-BEE5-EEC57FF2D024>

Figs 9, 10

Type material. *Holotype* ♂ (IZCAS-Ar40621), CHINA: Yunnan: Xishuangbanna, Mengla County, Menglun Town, Menglun Nature Reserve, Xishuangbanna Tropical Botanical Garden, tropical rainforest (21°55.35'N, 101°16.36'E, ca 610 m), 29.07.2018, X. Mi et al. leg. *Paratype* 1♀ (IZCAS-Ar40622), Palm Garden (21°55.47'N, 101°15.05'E, ca 550 m), 6.08.2018, C. Wang et al. leg.

Etymology. The specific name is a patronym after Prof. Xiaoqi Mi (Tongren, China), one of the collectors of the new species; noun (name) in genitive case.

Diagnosis. The male of *Phintella mii* sp. nov. resembles *P. aequipeiformis* Żabka, 1985 in the shape of the palp but can be distinguished by the following: 1) the distance between the embolus and cymbial tip is almost equal to the embolic length vs more than three times the embolic length in *P. aequipeiformis* (Luong et al. 2016: fig. 4A, D, E); 2) proximally, the RTA is strongly curved dorsally in retrolateral view vs straight in *P. aequipeiformis* (Luong et al. 2016: fig. 4B); 3) the chelicerae of the male are underdeveloped, the ratio of the length of the fang to the width of the paturon is about 1:1 vs well-developed chelicerae, with a ratio of almost 2.5:1 in *P. aequipeiformis* (Luong et al. 2016: fig. 3I). The female of this new species can be easily distinguished from other congeners by having an anterior subtriangular lamellar epigynal fold (Fig. 10A, F), which is absent in other species.

Description. Male (Figs 9, 10C, D, F, G). Total length 2.39. Carapace 1.05 long, 0.95 wide. Abdomen 1.15 long, 0.76 wide. Clypeus 0.05 high. Eye sizes and inter-distances: AME 0.30, ALE 0.15, PLE 0.14, AERW 0.93, PERW 0.91, EFL 0.59. Legs: I 2.46 (0.73, 0.90, 0.51, 0.32), II 2.20 (0.68, 0.76, 0.44, 0.32), III 2.68 (0.80, 0.88, 0.66, 0.34), IV 2.92 (0.90, 0.95, 0.73, 0.34). Carapace nearly square, bearing sparse white scales on cheeks, between both ALEs and PLEs and posteriorly, with a pair of indistinct stripes anteriorly and two pairs of round, dark markings near the PME and the posterior margin. Clypeus yellow-brown. Fovea longitudinal, short, bar-shaped. Chelicerae yellow, with one retromarginal tooth and two promarginal teeth. Endites yellow to green-brown. Labium dark brown. Sternum green-brown, heart-shaped, covered by dense, white setae. Legs pale to green-brown. Abdomen suboval, dorsum green-brown, speckled, with a nearly square white marking posteriorly, followed by a round, dark spot; venter dark brown, with a pair of longitudinal dotted lines medially.

Palp (Fig. 9A–D): tibia wider than long, with a RTA proximally curved dorsally, and then curved towards bulb terminally, pointed at the tip in retrolateral view; bulb flat, with sperm duct strongly bent anteriorly; lamellar process small, nearly round, regular bump located retrolatero-medially; embolus short, tapered, with a blunt tip directed antero-prolaterally in ventral view.

Female (Fig. 10A, B, E). Total length 3.04. Carapace 1.24 long, 1.01 wide. Abdomen 1.52 long, 1.12 wide. Clypeus 0.06 high. Eye sizes and inter-distances: AME 0.32, ALE 0.16, PLE 0.15, AERW 1.01, PERW 1.00, EFL 0.67. Legs: I 2.16 (0.68,




Figure 9. Male palp of the holotype of *Phintella mii* sp. nov. **A** prolateral **B** retrolateral **C** ventral **D** dorsal. Scale bars: 0.1.




Figure 10. *Phintella mii* sp. nov., female paratype and male holotype **A** epigyne, ventral **B** vulva, dorsal **C** holotype habitus, dorsal **D** holotype habitus, ventral **E** paratype habitus, dorsal **F** holotype carapace, dorsal **G** holotype chelicerae, frontal. Scale bars: 0.1 (**A**, **B**, **G**); 0.5 (**C**–**F**).

0.80, 0.44, 0.24), II 2.14 (0.68, 0.76, 0.46, 0.24), III missing, IV 3.05 (0.93, 1.05, 0.78, 0.29). Carapace yellow, with a pair of longitudinal dark stripes anteriorly in eye field and two pairs of dark brown patches close to PME and the posterior margin of the thoracic area. Abdomen suboval, dorsum pale, with indistinct transverse brown stripes medially and a round, brown spot at the terminus; venter colored as dorsum.

Epigyne (Fig. 10A, B): with a slightly recurved basal plate; copulatory openings located anteriorly, obscured by an inverted subtriangular lamellar epigynal fold; copulatory ducts extending transversely before extending posteriorly along the longitudinal axis, and then curving to connected to the dorsoectal surface of spermathecae; spermathecae almost round, separated from each other by less than one-third their diameter; fertilization ducts well-developed, located anteriorly on spermathecae.

Distribution. Known only from the type locality in Yunnan, China.

Comments. The female is paired with the holotype because it shares similar carapace markings with the male. These markings differ from other congeners known only from a single male from Xishuangbanna. However, the male and female have different abdominal markings, so the pairing requires further confirmation.

Genus *Simaetha* Strand, 1932

Type species. *Simaetha thoracica* Thorell, 1881

Comments. *Simaetha* is represented by 19 nominal species distributed in East and Southeast Asia, New Guinea, and Australia (WSC 2020). All species are endemic except *S. knowlesi* Żabka, 1994, *S. paetula* (Keyserling, 1882), *S. robustior* (Keyserling, 1882), *S. tenuidens* (Keyserling, 1882), and *S. tenuior* (Keyserling, 1882), known from both New Guinea and Australia. Although a comprehensive work was done by Żabka (1994), the genus is rather poorly studied, with 11 of its species known only from a single sex – five from females and six from males – and four species lack diagnostic drawings. To date, only the endemic species, *S. gongi* Peng, Gong & Kim, 2000, has been recorded from China.

Simaetha menglun sp. nov.

<http://zoobank.org/594A88B2-52FD-41BA-B066-97AB35F12359>

Figs 11, 12

Type material. *Holotype* ♂ (IZCAS-Ar40623), CHINA: Yunnan: Xishuangbanna, Mengla County, Menglun Town, Menglun Nature Reserve, Leprosy Village (21°53.62'N, 101°18.25'E, ca 520 m), 29.04.2019, Y. Tong et al. leg. *Paratype* 1 ♀ (IZCAS-Ar40624), same data as holotype.

Etymology. The species name is derived from the name of the type locality; noun in apposition.

Diagnosis. *Simaetha menglun* sp. nov. resembles *S. pengi* sp. nov. in the shape of the copulatory organs but can be easily distinguished by the shape of the habi-




Figure 11. Male palp of the holotype of *Simaetha menglun* sp. nov. **A** prolateral **B** retrolateral **C** ventral **D** bulb, ventral **E** dorsal. Scale bars: 0.1.

tus, the color pattern, and the cheliceral dentition. *S. menglun* sp. nov. resembles *S. gongji* Peng, Gong & Kim, 2000 by the general shape of the copulatory organs and the short palpal tibia but can be distinguished by the following: 1) the tip of

the embolus is curved in ventral view vs straight in *S. gongi* (Peng et al. 2000: fig. 14); 2) the epigynal hood is about one and a half times longer than wide vs almost two times wider than long in *S. gongi* (Peng et al. 2000: fig. 11); 3) the chelicerae of the female have one bifurcated retromarginal tooth vs two retromarginal teeth in *S. gongi* (Peng et al. 2000: fig. 10). The male of *S. menglun* sp. nov. also somewhat resembles *S. deelemanae* Zhang, Song & Li, 2003 in the shape of the palp, but can be easily distinguished by the palpal tibia which is wider than long rather than distinctly longer than wide in *S. deelemanae* (Zhang et al. 2003: fig. 7C, D).

Description. Male (Figs 11, 12D, E, G, H). Total length 2.49. Carapace 1.07 long, 1.01 wide. Abdomen 1.24 long, 0.88 wide. Clypeus 0.01 high. Eye sizes and inter-distances: AME 0.26, ALE 0.13, PLE 0.13, AERW 0.89, PERW 0.98, EFL 0.61. Legs: I 1.95 (0.68, 0.76, 0.29, 0.22), II 1.49 (0.49, 0.54, 0.24, 0.22), III 1.34 (0.44, 0.44, 0.24, 0.22), IV 1.79 (0.59, 0.61, 0.32, 0.27). Carapace nearly square, red-brown, with an irregular dark patch in the middle of the cephalic part, bearing dense, white setae on the cheeks, and dense, white and yellow setae dorsally. Clypeus dark. Fovea indistinct. Chelicerae red-brown, with two promarginal teeth and one retromarginal tooth bifurcated into two pointed tips. Endites and labium dark brown. Sternum colored as labium, covered with thin setae. Legs yellow to yellow-brown, with dark rings; legs I with dark, inflated tibia bearing dense setae ventrally. Abdomen elongated, dorsum with longitudinal, irregular, dark-brown stripe extending posteriorly from the anterior margin, followed by two chevrons, covered by dense, thin, white setae laterally; venter dark brown, with a pair of longitudinal white bands laterally.

Palp (Fig. 11A–E): tibia wider than long, with white scales dorsally and a lamellar RTA that tapers to a pointed tip which is almost directed towards 1:30 o'clock in retrolateral view; cymbium hirsute, with a proximo-retrolateral triangular process near the RTA base; bulb round and flat; embolus originates from a plate-like base, superimposed on the surface of the bulb, apically curved towards the prolateral side.

Female (Fig. 12A–C, F, I). Total length 3.01. Carapace 1.28 long, 1.09 wide. Abdomen 1.66 long, 1.23 wide. Clypeus 0.01 high. Eye sizes and inter-distances: AME 0.29, ALE 0.14, PLE 0.14, AERW 0.97, PERW 1.09, EFL 0.69. Legs: I 1.95 (0.66, 0.76, 0.29, 0.24), II 1.63 (0.51, 0.61, 0.27, 0.24), III 1.53 (0.49, 0.51, 0.29, 0.24), IV 2.03 (0.66, 0.76, 0.34, 0.27). Habitus similar to that of male except paler, and the chelicerae have three promarginal teeth.

Epigyne (Fig. 12A–C): wider than long, with an anterior hood approximately 1.5 times longer than wide; copulatory openings slit-like, located medially; copulatory ducts short, connected to the anterior chambers of the spermathecae; spermathecae divided into two round chambers; fertilization ducts connected to the anterior part of the posterior chambers of the spermathecae, extending obliquely.

Distribution. Known only from the type locality in Yunnan, China.

Comments. This species and *Simaetha pengi* sp. nov. are placed into this genus due to sharing a similar habitus and similar copulatory organs with *S. gongi* and *S. deelemanae* which occur in China and Singapore.




Figure 12. *Simaetha menglun* sp. nov., female paratype and male holotype **A, B** epigyne, ventral **C** vulva, dorsal **D** holotype habitus, dorsal **E** holotype habitus, ventral **F** paratype habitus, dorsal **G** holotype carapace, frontal **H** holotype chelicerae, dorsal **I** paratype chelicerae, dorsal. Scale bars: 0.1 (**A–C, H–I**); 0.5 (**D–G**).

***Simaetha pengi* sp. nov.**

<http://zoobank.org/DFE073DC-F938-4BEF-8492-B35D8716F8BA>

Figs 13, 14

Type material. *Holotype* ♂ (IZCAS-Ar40625), CHINA: Yunnan: Xishuangbanna, Mengla County, Menglun Town, Menglun Nature Reserve, Leprosy Village (21°53.62'N, 101°18.25'E, ca 520 m), 29.04.2019, Y. Tong et al. leg. *Paratypes* 2♀ (IZCAS-Ar40626–40627), same data as holotype.

Etymology. The specific name is a patronym in honor of Prof. Xianjin Peng (Changsha, China), who has produced many important taxonomic works on Chinese jumping spiders; noun (name) in genitive case.

Diagnosis. *Simaetha pengi* sp. nov. resembles *S. gongi* Peng, Gong & Kim, 2000 but can be distinguished by the apically curved embolus (vs straight), the longer than wide epigynal hood (vs wide than long), and the bifurcated retromarginal tooth of the female chelicerae (vs not bifurcated). The male of *S. menglun* sp. nov. also somewhat resembles *S. deelemanae* Zhang, Song & Li, 2003 in the shape of the palp but can be easily distinguished by the short palpal tibia, which is wider than long, but distinctly longer than wide in *S. deelemanae*.

Description. Male (Figs 13, 14C, D, F, G). Total length 2.17. Carapace 1.15 long, 1.11 wide. Abdomen 1.11 long, 1.01 wide. Clypeus 0.02 high. Eye sizes and inter-distances: AME 0.28, ALE 0.14, PLE 0.13, AERW 0.93, PERW 0.97, EFL 0.62. Legs: I 2.39 (0.78, 1.00, 0.34, 0.27), II 1.80 (0.56, 0.63, 0.34, 0.27), III 1.66 (0.51, 0.54, 0.34, 0.27), IV 2.00 (0.68, 0.71, 0.32, 0.29). Carapace almost square, red-brown, with a pair of indistinct dark brown patches medially on cephalic part, bearing dense, white setae forming two parallel transverse stripes on cheeks and golden-brown setae, two clusters of white setae near the PME on the dorsum. Clypeus dark brown. Fovea indistinct. Chelicerae red-brown, with two promarginal teeth and one retromarginal, pillar-shaped tooth bifurcated distally, with additional digitiform and round protuberances at the promargin and retromargin, respectively. Endites and labium colored as chelicerae. Sternum darker than endites, covered with pale, thin setae. Legs I with inflated, dark femora and tibiae, the latter bearing dense ventral setae, patellae yellow, tarsus pale yellow; other legs yellow with dark rings, except femora dark brown. Abdomen oval, dorsum dark medially, with spots of yellow scales anterolaterally and a yellow longitudinal band of scales posteriorly; venter brown.

Palp (Fig. 13A–D): tibia almost as long as wide, with transparent white scales dorsally, and a RTA less than tibial length, tapered to a slightly pointed tip, directed towards 1:00 o'clock in retrolateral view; cymbium hirsute, with a proximo-retrolateral triangular process and sparse, transparent white scales proximo-dorsally; bulb flat, with sperm duct extending along the margin; embolus arises from a plate-like base, superimposed on the surface of the bulb, slightly curved apically.

Female (Fig. 14A, B, E, H). Total length 2.48. Carapace 1.10 long, 1.03 wide. Abdomen 1.40 long, 1.13 wide. Clypeus 0.02 high. Eye sizes and inter-distances: AME 0.26, ALE 0.12, PLE 0.12, AERW 0.86, PERW 1.01, EFL 0.59. Legs: I 1.98 (0.71,




Figure 13. Male palp of the holotype of *Simaetha pengi* sp. nov. **A** prolateral **B** retrolateral **C** ventral **D** dorsal. Scale bars: 0.1.




Figure 14. *Simaetha pengi* sp. nov., female paratype and male holotype **A** epigyne, ventral **B** vulva, dorsal **C** holotype habitus, dorsal **D** holotype habitus, ventral **E** paratype habitus, dorsal **F** holotype carapace, frontal **G** holotype chelicerae, dorsal **H** paratype chelicerae, dorsal. Scale bars: 0.1 (**A, B, G, H**); 0.5 (**C-F**).

0.76, 0.27, 0.24), II 1.64 (0.54, 0.59, 0.27, 0.24), III 1.53 (0.49, 0.51, 0.29, 0.24), IV 2.03 (0.71, 0.73, 0.32, 0.27). Habitus similar to that of male, except paler, and the chelicerae have one promarginal tooth and one retromarginal fissident with three cusps.

Epigyne (Fig. 14A, B): wider than long, with an anterior bell-shaped hood approximately 1.5 times longer than wide; copulatory openings slit-like, located medially, separated from each other by approximately 1.5 times the width of the hood; copulatory ducts short; spermathecae divided into two round chambers; fertilization ducts connected to the anterior part of the posterior chambers of the spermathecae.

Distribution. Known only from the type locality in Yunnan, China.

Acknowledgements

The manuscript benefited greatly from comments by Yuri Marusik (Magadan, Russia), Dmitri Logunov (Manchester, UK), Galina N. Azarkina (Novosibirsk, Russia), Wayne Maddison (Vancouver, Canada), and an anonymous referee. Łukasz Trębicki confirmed the identification of *Charippus yinae* sp. nov. Sarah Crews (San Francisco, USA) checked the English of the final manuscript. Yanfeng Tong, Hao Yu, Zhigang Chen, Zilong Bai, Shijia Liu, Xiaoqi Mi, Jiahui Gan, Yuanfa Yang, and Hong Liu helped with fieldwork. This research was supported by the National Natural Science Foundation of China to Shuqiang Li (NSFC-31530067), Xiaoqi Mi (NSFC-31660609), and Natural Science Research Project Foundation of Guizhou Province of Education (KY-2018-345).

References

- Benjamin SP (2004) Taxonomic revision and phylogenetic hypothesis for the jumping spider subfamily Ballinae (Araneae, Salticidae). *Zoological Journal of the Linnean Society* 142(1): 1–82. <https://doi.org/10.1111/j.1096-3642.2004.00123.x>
- Cao Q, Li S, Žabka M (2016) The jumping spiders from Xishuangbanna, Yunnan, China (Araneae, Salticidae). *ZooKeys* 630: 43–104. <https://doi.org/10.3897/zookeys.630.8466>
- Davies VT, Žabka M (1989) Illustrated keys to the genera of jumping spiders (Araneae: Salticidae) in Australia. *Memoirs of the Queensland Museum* 27: 189–266.
- Li S (2020) Spider taxonomy for an advanced China. *Zoological Systematics* 45(2): 73–77.
- Lin Y, Li S (2020) Two new genera and eight new species of jumping spiders (Araneae, Salticidae) from Xishuangbanna, Yunnan, China. *ZooKeys* 952: 95–128. <https://doi.org/10.3897/zookeys.952.51849>
- Luong PTH, Yamasaki T, Eguchi K (2016) Conspecificity of *Phintella aequipeiformis* Zabka, 1985 and *P. lucai* Zabka, 1985 (Araneae: Salticidae) confirmed by DNA barcoding. *Revue Suisse de Zoologie* 123(2): 283–290.
- Malamel JJ, Prajapati DA, Sudhikumar AV, Sebastian PA (2019) Two new species of the tribe Ballini Banks, 1892 from India (Araneae: Salticidae). *Arthropoda Selecta* 28(3): 424–434. <https://doi.org/10.15298/arthsel.28.3.07>

- Myers N (1988) Threatened biotas: “hot spots” in tropical forests. *The Environmentalist* 8: 1–20. <https://doi.org/10.1007/BF02240252>
- Peng XJ (1995) Two new species of jumping spiders from China (Araneae: Salticidae). *Acta Zootaxonomica Sinica* 20: 35–38.
- Peng XJ, Gong LS, Kim JP (2000) Two new species and two unrecorded species of the family Salticidae (Arachnida: Araneae) from China. *Korean Journal of Soil Zoology* 5: 13–19.
- Peng XJ, Kim JP (1997) Three new species of the genus *Eupoa* from China (Araneae: Salticidae). *Korean Journal of Systematic Zoology* 13: 193–198.
- Peng XJ, Xie LP, Xiao XQ, Yin CM (1993) Salticids in China (Arachnida: Araneae). Hunan Normal University Press, 270 pp.
- Peng XJ, Yin CM (1991) Five new species of the genus *Kinbia* from China (Araneae: Salticidae). *Acta Zootaxonomica Sinica* 16: 35–47.
- Song DX (1991) Three new species of the genus *Ptocasius* from China (Araneae: Salticidae). *Sinozoologia* 8: 163–168.
- Song DX, Zhu MS (1998) Two new species of the family Salticidae (Araneae) from China. *Acta Arachnologica Sinica* 7: 26–29.
- Thorell T (1895) *Descriptive Catalogue of the Spiders of Burma, Based upon the Collection Made by Eugene W. Oates and Preserved in the British Museum*. London, 406 pp.
- Wang C, Li S (2020) On eight species of jumping spiders from Xishuangbanna, Yunnan, China (Araneae, Salticidae). *ZooKeys* 909: 25–57. <https://doi.org/10.3897/zookeys.909.47137>
- Wanless FR (1988) A revision of the spider group Astieae (Araneae: Salticidae) in the Australian region. *New Zealand Journal of Zoology* 15: 81–172. <https://doi.org/10.1080/03014223.1988.10422611>
- WSC (2020) World Spider Catalog, version 21.5. Natural History Museum Bern. <http://wsc.nmbe.ch> [2020-8-23]
- Xiao XQ (2002) A new species of the genus *Myrmarachne* from China (Araneae: Salticidae). *Acta Zootaxonomica Sinica* 27: 477–478.
- Xiao XQ, Wang SP (2004) Description of the genus *Myrmarachne* from Yunnan, China (Araneae, Salticidae). *Acta Zootaxonomica Sinica* 29: 263–265.
- Xie LP, Peng XJ (1995) Spiders of the genus *Thyene* Simon (Araneae: Salticidae) from China. *Bulletin of the British Arachnological Society* 10: 104–108.
- Żabka M (1985) Systematic and zoogeographic study on the family Salticidae (Araneae) from Viet-Nam. *Annales Zoologici, Warszawa* 39: 197–485.
- Żabka M (1992) Salticidae (Arachnida: Araneae) from Oriental, Australian and Pacific regions, VII. *Paraplatooides* and *Grayenulla* – new genera from Australia and New Caledonia. *Records of the Australian Museum* 44: 165–183. <https://doi.org/10.3853/j.0067-1975.44.1992.31>
- Żabka M (1994) Salticidae (Arachnida: Araneae) of Oriental, Australian and Pacific regions, X. Genus *Simaetha* Thorell. *Records of the Western Australian Museum* 16: 499–534.
- Zhang JX, Song DX, Li DQ (2003) Six new and one newly recorded species of Salticidae (Arachnida: Araneae) from Singapore and Malaysia. *Raffles Bulletin of Zoology* 51: 187–195.