

# A gynandromorphy of *Polistes nimphus* (Christ, 1791) (Hymenoptera: Vespidae: Polistinae) in China

YUAN Feng<sup>1</sup>, ZHAO Yan-Hui<sup>1,2</sup>, ZHANG Yan-Zhou<sup>1</sup>, ZHU Chao-Dong<sup>1,\*</sup>

(1. Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China)

2 College of Life Science, Capital Normal University, Beijing 100048, China

**Abstract** This note reports the occurrence of gynandromorphism in *Polistes nimphus* (Christ 1791) in China. External features of the insect are presented. The insect was collected in Mt Donglingshan, Mentougou, Beijing. The specimen studied is deposited in the Institute of Zoology, Chinese Academy of Sciences.

**Key words** Hymenoptera Vespidae *Polistes nimphus* morphological characters gynandromorphism

Within the family Vespidae (Hymenoptera), gynandromorphism has been recorded in 3 subfamilies 3 genera 2 species of which are in the genus *Polistes* Latreille, 1802 (Table 1). The mechanism of this phenomenon is not completely known, but it is generally attributed to developmental anomalies. In addition, low temperature might induce gynandromorphism (Wilson, 1962; Cui and Cai 2003; Zhang and Zhu 2007).

**Table 1** Gynandromorphism in Vespidae

Family/species	References
Vespinae	
<i>Vespa crabro</i> Linnaeus 1758	Dessart 1990
Polistinae	
<i>Polistes gallicus</i> (Linnaeus 1767)	Gauss 1966
<i>Polistes dominulus</i> (Christ 1791)	Wolf 1985
Eumeninae	
<i>Odynerus reniformis</i> (Gmelin 1790)	Leininger 1926

The genus *Polistes* was established by Latreille (1802). The type species of the genus is *Vespa gallica* Linnaeus 1767. *Polistes* Latreille has been divided into four subgenera, *Aphanilopterus*, *Gyrostoma*, *Polistella* and *Polistes* s. str., with 212 species in the world (fossil species and unrecognized ones). Twenty-three species of the genus *Polistes* have been reported in China, of which 5 species are in the subgenus *Polistes* s. str. (Carpenter 1996). They are *P. associus* Kohl 1898, *P. chinensis* (Fabricius

1793), *P. dominulus* (Christ 1791), *P. gallicus* (Linnaeus 1767) and *P. nimphus* (Christ 1791).

In China, 3 species are catalogued by Liu (1937) in the subgenus *Polistes* s. str. They are *P. chinensis* (Fabricius 1793), *P. dominulus* (Christ 1791) and *P. gallicus* (Linnaeus 1767). Two more species were reported by Li (1982, 1985), which are *P. chinensis* (Fabricius 1793), *P. gallicus* (Linnaeus 1767). Li (2006) moved specimens under the name *P. gallicus* (Linnaeus 1767) to *P. dominulus* (Christ 1791), without providing any evidences.

On 28th Sept 2008, one gynandromorphous adult specimen of the genus *Polistes* Latreille was collected by YUAN Feng and ZHAO Yan-Hui with yellow pan-trap in Mt Donglingshan ( $40^{\circ}04'N$ ,  $115^{\circ}49'E$ , elevation 1 154 m), Mentougou, Beijing, China. Examined related contents especially digital images from [http://userswing.be/entomobie/Vespidae\\_2.htm](http://userswing.be/entomobie/Vespidae_2.htm) (Reneson 2008), we identified this gynandromorphous adult as *P. nimphus* (Christ 1791). Since little has been known about *P. nimphus* on its morphological anomalies, here we describe the external features related to the gynandromorphism in *P. nimphus* (Christ 1791). In both dorsal and ventral view, the body of the gynandromorphous adult is divided symmetrically in length, with male external features on the right and female ones on the left (Table 2, Figs 4–5). Sexual dimorphism presents its antennae, head front and lateral parts of the body (shape, size and color) (Figs 1, 2, 3, 6). There are both male and female features of genitalia (Figs 7–8).

基金项目：中国农业部行业科技专项（200803006）


作者简介：袁峰，男，1974年4月生，北京人，从事昆虫分类学研究，Tel 13693017579，E-mail yuan@ioz.ac.cn

\* 通讯作者 Author for correspondence E-mail zhuced@ioz.ac.cn

收稿日期 Received 2009-04-13 接受日期 Accepted 2009-07-08

© 1994-2011 China Academic Journal Electronic Publishing House. All rights reserved. <http://www.cnki.net>

2003-1-1, 0:08

Figs 1–8 Gynandromorph of *Polistes nimphus* (Christ 1791)

1. Antennae and head, dorsal view; 2. Head, front view; 3. Body, lateral view; 4. Body, dorsal view; 5. Body, ventral view; 6. Body, right lateral view; 7. Apex of abdomen, dorsal view; 8. Apex of abdomen, ventral view.

Table 2 A comparison of main features of gynandromorphy

Body parts	Female (left)	Male (right)
Head	Face at top of antecava socket inner margin of eye end of gena base and end edge of clypeus base of clypeus with yellow spots antennawith 10 flagellomeres	Face at the lower half of frons clypeus gena and mandible yellow; antennawith 11 flagellomeres
Thorax	Mesopleural epipleurite with yellow spots	Mesopleural epipleurite and hypopleurite with yellow spots
Abdomen	Abdomen with six abdominal gastral tergum II with yellow spots small gastral sternite II – III without yellow spots	Abdomen with seven abdominal gastral tergum II with yellow spots large gastral sternite II – III with yellow spots

**ACKNOWLEDGEMENTS** We thank Mr LIU Ye in the Institute of Zoology Chinese Academy of Sciences for taking photos of the adult specimen. This project was supported mainly by the Innovation Program in the Chinese Academy of Sciences and partially by Public Welfare Project from the Ministry of Agriculture China (Grant No. 200803006).

## References

- Carpenter JM, 1996. Distributional checklist of species of the genus *Polistes* (Hymenoptera: Vespidae; Polistinae; Polistini). *American Museum Novitates*, 3188: 1–39.
- Christ JL, 1791. *Naturgeschichte, Classification und Nomenclatur der Insecten von Bienen, Wespen, und Ameisengeschlecht als der fuenften Klasse fuenften Ordnung des Linneischen NaturSystems von den Insecten Hymenoptera*. Hemann, Frankfurt am Main 1–535.
- Cui JX, Cai WZ, 2003. Gynandromorphism in insects. *Entomological Knowledge*, 40(6): 565–570. [崔建新, 彩万志, 2003. 昆虫的雌性嵌合现象. 昆虫知识, 40(6): 565–570]
- Dessart P, 1990. Un exemple de frelon (*Vespa crabro* Linne 1758). *Bulletin et Annales de la Societe Royale Belge d'Entomologie*, 126 (10/12): 206.
- Gauss R, 1966. Cyclopie bei *Pleolophus brachypterus* (Grau) (Hym. Ichneumon), Gynander von *Polistes gallicus* (Linne) (Hym. Vespidae) und Kopulations-Anomalie von *Melolontha melolontha* (Linne) (Col. Lamellae). *Deutsche Entomologische Zeitschrift*, 13: 291–294.
- Leininger H, 1926. Ein lateraler Zwitter von *Odynerus reniformis* Gmel (Hym., Vespidae). *Verh. naturw. Ver. Karlsruhe*, 30: 162–167.
- Li TS, 1982. Homets from Agriculture of China (Hymenoptera Vespoidea). Agriculture Press, Beijing 255 pp. [李铁生, 1982 中国农区胡蜂. 北京: 农业出版社. 255页]
- Li TS, 1985. Economic Insect Fauna of China. Fasc. 30, Hymenoptera Vespoidea. Science Press, Beijing 159 pp. [李铁生, 1985 中国经济昆虫志, 第三十册 膜翅目: 胡蜂总科. 北京: 科学出版社. 159页]
- Li TS, 2006. Secrets of Wasps. China Forestry Publishing House, Beijing 72 pp. [李铁生, 2006 马蜂揭秘. 北京: 中国林业出版社. 72页]
- Linnaeus C, 1767. *Systema naturae per regna tria naturae secundum classes, ordines, genera, species cum characteribus differentiis, synonymis, 12th ed. Laur. Silv. Holmiae*.
- Liu CL, 1937. A bibliographic and synoptic catalogue of the Vespidae of China with cross-reference index for the genera and species. *Peking Natural History Bulletin*, 11(3): 205–232.
- Renneson JL, 2008. Hymenoptera Vespidae. [http://users.swing.be/entomologie/Vespidae\\_2.htm](http://users.swing.be/entomologie/Vespidae_2.htm). [Updated October 27]
- Wilson F, 1962. Sex determination and gynandromorph production in aberrant and normal strains of *Ooencyrtus submetallicus* (Hymenoptera Encyridae). *Australian Journal of Zoology*, 10 (3): 349–359.
- Wolf H, 1985. Zwitter von *Evaetus crassicornis* (Shuckard) (Hym., Pompilidae), *Polistes dominulus* (Christ) (Hym., Vespidae), *Crabro peltarius* (Schreber) und *Ceræris arenaria* (Linne) (Hym., Sphecidae). *Linzer Biologische Beiträge*, 17(2): 495–498.
- Zhang YZ, Zhu CD, 2007. A gynandromorph of *Microterys ishii* Tadikawa (Hymenoptera Chalcidoidea Encyridae). *Acta Entomologica Sinica*, 50(8): 868–870. [张彦周, 朱朝东, 2007 石井花跳小蜂(膜翅目: 跳小蜂科)的雌雄同体个例. 昆虫学报, 50(8): 868–870]

## 普通马蜂的雌雄嵌体个例

袁峰<sup>1</sup>, 赵延会<sup>1,2</sup>, 张彦周<sup>1</sup>, 朱朝东<sup>1,\*</sup>

(1. 中国科学院动物研究所动物进化与系统学重点实验室, 北京 100101; 2. 首都师范大学生命科学学院, 北京 100048)

**摘要:** 本文报道普通马蜂 *Polistes nimphus* (Christ 1791) 的雌雄嵌体一例, 并提供其形态特征图。研究标本的采集地为北京门头沟区东灵山, 现保存在中国科学院动物研究所动物标本馆。

**关键词:** 膜翅目; 胡蜂科; 普通马蜂; 形态特征; 雌雄嵌体

**中图分类号:** Q969   **文献标识码:** A   **文章编号:** 0454-6296(2009)09-1055-03

(责任编辑: 袁德成)